用藥指導單張

衛教小組 DJ13580101

學　名：Zn+Cu+Mn+Se+F+I

商品名：PEDITRACE INJ. 10ML

中文名：



一、此藥品的用途是什麼？

To cover basal trace element needs for neonates and children receiving total intravenous nutrition.

二、此藥品該如何使用？

詳細的使用方式請遵從藥袋上用法用量醫師的指示。一般來說這個藥品的使用方式如下：

Must be diluted.

1 ml Peditrace per kg of body weight per day for infants and children weighing up to 15 kg. For children weighing 15 kg or more, a daily dose of 15 ml Peditrace covers the basal need of trace elements. The infusion duration should be at least 8 hours.

三、使用時我該注意的特別事項？

Peditrace should be used with caution in patients with impaired biliary and/or renal function, in whom the excretion of trace elements may be significantly decreased.

Peditrace should also be used with caution in patients with biochemical or clinical evidence of liver dysfunction (especially cholestasis). Trace element levels in plasma must be monitored in patients with cholestatic liver disease and renal insufficiency. If treatment lasts more than 4 weeks, manganese levels must be checked.

If intravenous nutrition is administered for a prolonged period of time (month), plasma levels of trace elements should also be monitored.

四、孕婦用藥等級及注意事項？

請諮詢醫師或藥師

五、此藥會造成什麼副作用？

No side effects related to the trace elements in Peditrace have been reported.

六、此藥儲存時的條件為何？

Store at a maximum of 25° C. Do not freeze.

本單張資料僅供參考，如有疑問請諮詢藥師

光田醫療社團法人 光田綜合醫院藥劑部
諮詢電話　向上/沙鹿院區 (04)26625111-1050
大甲院區 (04)26885599-5141
審視日期：2026.11.16